Mepore Film How To

How To Use

Mepore® Film

Transparent, breathable, adhesive coated polyurethane film

Product description

Mepore Film is a flexible, transparent, breathable film dressing made of polyurethane, coated with a polyacrylic adhesive. The dressing is supported with a transparent sheet paper frame to ease the application.

Mode of action

Mepore Film provides a flexible, transparent covering to protect the wound whilst conforming to surface irregularities and body contours.

Mepore Film maintains a moist environment yet due to its vapour permeability allows excess moisture pass away from the wound and surrounding skin.

Mepore Film provides a barrier to fluid strike-through and bacterial contamination.

Intended use

Mepore Film is intended for a wide range of wounds in the granulation phase such as superficial burns, IV sites, abrasions, lacerations, donor sites, superficial pressure ulcers, closed surgical wounds and may help to prevent skin breakdown.

Mepore Film can be used as a conformable fixation device or as a secondary dressing to cover dressings such as hydrogels, alginates and foam dressings.

Precautions

Mepore Film should not be applied on patients who are sensitive to acrylic adhesive.

Mepore Film should not be used directly on full thickness wounds involving muscle, tendon or bone or on third degree burns.

Method of use

Application:

- 1. Clean the application area. Dry the healthy surrounding skin.
- 2.Chose the correct dressing size to allow a margin which adheres to dry healthy skin around the wound or catheter site. The dressing may be cut and can be overlapped.
- 3. For size 10x25 cm (4x10 inches) and 15x20 cm (6x8 inches) only: Remove center cut-out and discard.
- 4. Remove the protection paper (printed Mepore) to expose the adhesive.
- 5. Apply Mepore Film and remove the paper frame.
- 6. Remove the two white paper side tabs.

Dressing change and removal

- 1. The dressing may be left in place for several days depending on the wound/skin condition, or as indicated by good clinical practice.
- 2. Lift a corner of the dressing and gently pull the film along the skin (in the direction of hair growth).

Sterility is guaranteed unless the inner package is damaged or opened prior to use. Do not resterilise.